

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Biochemistry -- Faculty Publications

Biochemistry, Department of

1994

Characteristics of Modified Leghemoglobins Isolated from Soybean (*Glycine max* Merr.) Root Nodules

Hyung-Kyun Jun

University of Nebraska - Lincoln

Gautam Sarath

University of Nebraska-Lincoln, Gautam.sarath@ars.usda.gov

Jose F. Moran

Consejo Superior de Investigaciones Científicas

Manuel Becana

Consejo Superior de Investigaciones Científicas

Robert V. Klucas

University of Nebraska - Lincoln

See next page for additional authors

Follow this and additional works at: <http://digitalcommons.unl.edu/biochemfacpub>



Part of the [Biochemistry Commons](#), [Biotechnology Commons](#), and the [Other Biochemistry, Biophysics, and Structural Biology Commons](#)

Jun, Hyung-Kyun; Sarath, Gautam; Moran, Jose F.; Becana, Manuel; Klucas, Robert V.; and Wagner, Fred W., "Characteristics of Modified Leghemoglobins Isolated from Soybean (*Glycine max* Merr.) Root Nodules" (1994). *Biochemistry -- Faculty Publications*. 332. <http://digitalcommons.unl.edu/biochemfacpub/332>

This Article is brought to you for free and open access by the Biochemistry, Department of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Biochemistry -- Faculty Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Authors

Hyung-Kyun Jun, Gautam Sarath, Jose F. Moran, Manuel Becana, Robert V. Klucas, and Fred W. Wagner

Characteristics of Modified Leghemoglobins Isolated from Soybean (*Glycine max* Merr.) Root Nodules¹

Hyung-Kyun Jun, Gautam Sarath, Jose F. Moran, Manuel Becana, Robert V. Klucas, and Fred W. Wagner*

Department of Biochemistry, University of Nebraska-Lincoln, Lincoln, Nebraska 68583-0718 (H-K.J., G.S., R.V.K., F.W.W.); and Estación Experimental de Aula Dei, Consejo Superior de Investigaciones Científicas, Apartado 202, 50080 Zaragoza, Spain (J.F.M., M.B.)

Hemoprotein derivatives of an abundant soybean (*Glycine max* Merr.) root nodule leghemoglobin, Lba, were studied for their modified spectral characteristics and physical properties. Three modified hemoprotein derivatives of Lba (Lbam₁, Lbam₂, and Lbam₃) were purified by preparative isoelectric focusing. The ferric forms of these pigments were green and exhibited anomalous spectra in the visible region as compared to the Lba³⁺ forms. These modified pigments showed a hypochromic shift of 10 nm for the charge transfer absorption maximum; however, differences were not apparent in the Soret region. Upon binding with nicotinate, the α and β bands were shifted significantly into the red region as compared to the Lba³⁺ nicotinate complex. The three Lbam fractions were reduced by dithionite or by NADH in the presence of riboflavin. Lbam²⁺ also bound nicotinate and displayed absorption spectra indistinguishable from those of Lba²⁺ nicotinate. In contrast to Lba²⁺, Lbam²⁺ displayed aberrant spectra when bound with either O₂ or CO. These complexes exhibited a prominent charge transfer band at approximately 620 nm and failed to exhibit spectra characteristic of Lba²⁺O₂ and Lba²⁺CO. The protein moiety of these modified pigments was intact because their tyrosine/tryptophan ratios and their amino acid compositions were identical with those of Lba, nor were differences observed in the peptide profiles resulting from trypsin digests of purified Lba and Lbams. Automated Edman degradation of selected peaks further confirmed the intactness of the protein backbone including the absence of deamination. Pyridine hemochromogen for heme from Lbams could be formed, and the spectra displayed distinct differences compared to those of Lba. A new peak at 580 nm and a loss of a peak at 480 nm were observed for all three Lbams.

Lbs are monomeric hemoproteins that facilitate O₂ diffusion within legume root nodules (Appleby, 1984). Like other O₂-carrying hemoproteins, such as myoglobin and hemoglobin, Lbs display reversible oxidation and reduction of the heme iron. To bind O₂ reversibly, Lbs must be reduced to their ferrous forms. Whereas studies have shown that soybean (*Glycine max* Merr.) Lb levels decline with the onset of nodule senescence, whether induced or natural (Klucas, 1974;

Pfeiffer et al., 1983; Sutton, 1983), the biochemical events surrounding this decline are unknown. It is also unclear whether Lbs undergo irreversible changes during repeated oxidation-reduction cycles and whether inactive Lbs are present in appreciable quantities in functional root nodules.

An analysis of soybean Lbs by ion-exchange HPLC revealed the presence of a hemoprotein that appeared to be very similar to Lba but with somewhat different properties (Sarath et al., 1986). The ratio of the levels of this protein, termed Lbam (Wagner and Sarath, 1987), to Lba increased when nodulated soybean plants were subjected to dark stress. Lbam was the major hemoprotein detectable in nodules at advanced stages of senescence (Wagner and Sarath, 1987). Preliminary spectroscopic data indicated that in the ferric state Lbam was different from Lba, although this Lbam³⁺ preparation underwent normal reduction (Wagner and Sarath, 1987). These observations suggested that Lbam was a potential inactive form of Lba that accumulated in senescent nodules. However, the small amounts of Lbam, purified by HPLC, precluded a detailed investigation of its properties.

Recently, a method was developed to separate and purify Lbams by ion-exchange chromatography, followed by preparative IEF (Jun, 1993). This procedure results in the separation of a crude Lbam fraction into three distinct proteins resolvable by IEF. All three Lbams possessed an NH₂-terminal sequence identical with that of Lba and only appeared to differ in their isoelectric point (Jun, 1993), suggesting that subtle changes had occurred in these molecules. In this report, we demonstrate that these proteins exhibit characteristic spectral properties and show anomalous binding of O₂ and CO.

MATERIALS AND METHODS

Protein Purification

Root nodules were harvested from 40- to 50-d-old *Glycine max* Merr. cv Hobbit plants nodulated with *Bradyrhizobium japonicum* strain 61A89. Lbs were purified from soybean root nodules according to the method of Jun (1993). Briefly, root nodule extracts were prepared in 20 mM Na-phosphate buffer (pH 6.8) and fractionated with ammonium sulfate. Proteins

¹ This work was supported by grants from the Nebraska Soybean Development, Utilization and Marketing Board (F.W.W.), The Center for Biotechnology, University of Nebraska-Lincoln (G.S.), and Dirección General de Investigación Científica y Tecnológica (grant PB92-0058; M.B.). This is published as paper No. 10577, Journal Series, Agriculture Research Division, University of Nebraska-Lincoln.

* Corresponding author; fax 1-402-472-7842.

Abbreviations: Lb, leghemoglobin; Lb²⁺, ferrous leghemoglobin; Lb³⁺, ferric leghemoglobin; Lba, leghemoglobin component a; Lbam_x, modified components of Lba (x = 1, 2, or 3); Lbb, leghemoglobin component b.

precipitating between 35 and 90% saturation were dialyzed against 10 mM Na-phosphate (pH 6.8) and applied to an hydroxylapatite column equilibrated and washed with the same buffer. The column effluent containing Lbs was concentrated and dialyzed against 20 mM Tris-HCl (pH 9.2). Lb samples were subsequently oxidized with an excess of ferricyanide (Appleby et al., 1975) and chromatographed in a Sephadex G-75 column with 20 mM Tris-HCl (pH 9.2) containing 0.6 M NaCl. Fractions containing Lbs were concentrated, dialyzed against 20 mM Tris-HCl (pH 8.0), and applied to a DEAE-cellulose (Whatman DE-52) column. This column was developed isocratically with 20 mM Tris-HCl (pH 8.0). Two major Lb fractions, *Lba* and *Lbams*, were collected. Each of these fractions was further purified as Lb^{3+} nicotinate complexes by preparative IEF gels containing 2% of pH 4 to 6 ampholytes (Jun, 1993). The catholyte was 0.1 M β -Ala, and the anolyte was 0.1 M acetic acid. The *Lbam* fraction from the DEAE-cellulose column was resolved into three major proteins, namely *Lbam*₁, *Lbam*₂, and *Lbam*₃. The resolved proteins were eluted from macerated gel pieces with 20 mM Tris-HCl (pH 9.2). Pooled fractions were concentrated, oxidized with an excess of ferricyanide (Appleby et al., 1975), and passed over a PD-10 column (Pharmacia Fine Chemicals) equilibrated and developed with 20 mM Tris-HCl (pH 9.2). Recovered Lb^{3+} samples were concentrated and buffer exchanged into 50 mM K-phosphate buffer (pH 7.0) using a PD-10 column. Lbs were stored in a concentrated form (1–4 mM) at -20°C . All samples were concentrated using YM-10 Centricon concentrators (Amicon Corp.).

The concentrations of both *Lba* and *Lbams* possessing a single band in SDS-PAGE were determined by using the molar absorptivity of *Lba* at 280 nm ($29\text{ cm}^{-1}\text{ mM}^{-1}$) (Appleby et al., 1975).

Spectroscopic Methods

Spectra of purified proteins were performed in 50 mM K-phosphate buffer (pH 7.0). Visible spectra were collected using a Hitachi U-2000 spectrophotometer interfaced to a microcomputer or on a Cary 2200 spectrophotometer.

Lb^{3+} samples were either reduced with a few crystals of dithionite (Becana and Klucas, 1990) or with a mixture of riboflavin and NADH (Becana et al., 1991). Reduced Lb samples were mixed with O_2 -saturated or CO-saturated buffers to obtain these liganded forms. Pyridine-hemochromogen spectra were obtained as described earlier (Riggs, 1981).

Protein Analyses

Apo-Lbs were generated by extraction of acidified protein samples with 2-butanone (Ascoli et al., 1981). To avoid protein precipitation, apo-Lb samples were neutralized immediately to pH 7.0 by the addition of a 50- μL aliquot of 1 M K-phosphate (pH 7.0), followed by dialysis against 50 mM K-phosphate (pH 7.0) to remove all traces of 2-butanone. Concentrated apo-Lb samples were used for the determination of their Tyr/Trp ratios (Edelhoch, 1967).

Trypsin Digestion

N-tosyl-L-Phe chloromethyl ketone-treated trypsin (Sigma) was dissolved in 0.5 mM HCl. An 8- μg aliquot of trypsin was added to 400 μL of Lb sample (1 $\mu\text{g}/\mu\text{L}$) in 20 mM Tris-HCl (pH 8.0) containing 10 mM CaCl_2 . After an initial incubation for 2 h at 37°C , an additional 8- μg aliquot of trypsin was added and digestion was allowed to proceed for an additional 3 h. Enzymic reactions were terminated by the addition of 8 μL of TFA.

Released peptides were analyzed by HPLC (Sarath et al., 1986). Peptides were resolved on a $3.9 \times 150\text{-mm}$ C_{18} - $\mu\text{Bondapak}$ column (Waters Associates) at room temperature using gradients with acetonitrile. Multiple injections of the same samples were performed using an automated sample injection (WISP 715, Waters Associates), and the column eluate was analyzed at A_{215} and subsequently at A_{290} . Two peptides with high A_{290} were further analyzed by automated Edman degradation on a Millipore 6600 protein sequencer using protocols supplied by the manufacturer.

RESULTS AND DISCUSSION

Lb^{3+} Spectra

The absorption spectra for the *Lba*³⁺ and *Lbam*³⁺s are shown in Figure 1. Several features characteristic of the *Lbm*³⁺ were readily apparent. The charge transfer bands in the Lb^{3+} state were shifted between 13 and 20 nm into the blue region for the *Lbams*. In addition, the α band was pronounced between 551 and 555 nm, and the β band at 530 nm was absent. The *Lbm*³⁺s showed lower absorptivities in their Soret bands compared to *Lba*³⁺. The nicotinate complexes of *Lbm*³⁺s exhibited significantly different α and β bands. Both bands were shifted bathochromically over 20 nm in the α band and over 14 nm in the β band. *Lbam*₁³⁺ and *Lbam*₃³⁺ exhibited a 4-nm red shift in the Soret region compared to the spectra observed with *Lba*³⁺. These spectral differences could arise from changes to either the globin component or to the heme. The fact that nicotinate would bind to *Lbams* shows that the sixth coordination site on the iron is accessible to the solvent and, therefore, to ligands. However, the spectral shifts and the anomalous spectra of *Lbm*³⁺ indicate that some changes to the heme or heme environment had taken place that did not involve a new ligand.

Lb^{2+} Spectra and Ligand-Binding Properties

Spectra of the *Lbm*²⁺ and *Lbm*²⁺-nicotinate were very similar to those of *Lba* (Fig. 2), indicating that the heme in the *Lbm* molecules could undergo reduction and binding of nicotinate in a manner similar to those of *Lba*. Analyses of *Lbm*²⁺s in the presence of either O_2 or CO did not reveal the characteristic appearance of strong absorption bands at 574 and 540 nm of the *Lba*²⁺ O_2 complex or the 562- and 538-nm bands of the *Lba*²⁺CO complex (Fig. 3). Instead, the three *Lbams* displayed a strong charge transfer band at approximately 618 to 625 nm and minor bands centered at approximately 570 and 535 nm but failed to exhibit a clearly defined peak in the Soret region. Earlier data that indicated




Figure 1. Visible absorption spectra of Lb^{3+} and Lb^{3+} nicotinate complexes for *Lba* and *Lbam*s. Samples purified from preparative IEF gels were oxidized with a 4-fold excess of ferricyanide, desalted by gel filtration at pH 9.2, concentrated, and buffer exchanged in 50 mM K-phosphate (pH 7.0). Nicotinate complexes were formed by incubating Lb^{3+} samples with a final nicotinate concentration of 0.5 mM. Samples were scanned from 350 to 450 nm (8 μ M *Lb* in 500 μ L total volume) and from 450 to 700 nm (50 μ M *Lb* in 500 μ L total volume) with a Hitachi U-2000 spectrophotometer. The spectra were obtained after 20 min of equilibration at 22°C. Solid line, Lb^{3+} ; dotted line, Lb^{3+} nicotinate; A, *Lba*; B, *Lbam*₁; C, *Lbam*₂; D, *Lbam*₃.

CO binding with an *Lbam* fraction (Wagner and Sarath, 1987) can probably be ascribed to the presence of *Lba* in those *Lbam* samples. It is also possible that *Lbams* isolated from senescent root nodules might have slightly different properties.

The apparent aberrant binding of *Lbams* with O_2 could have important effects on nodule metabolism. If such proteins become predominant, and as our data suggest have impaired capacity to reversibly bind O_2 , the flux of O_2 to the bacteroids could decrease, which is in accord with studies showing an increase in the amounts of *Lbams* with the onset of stress (Wagner and Sarath, 1987).

Protein Structure Studies

The apparent changes in the binding of O_2 and CO suggested that a subtle modification of *Lba* had taken place to form the *Lbams*. To evaluate the causes for this change, a detailed examination of the globin backbone was undertaken.

Lehtovaara (1978) had suggested that a specific Tyr residue near the COOH terminus in bean *Lb* appeared to be essential in protecting the heme from hydroxylation reactions. The loss of this Tyr appeared to be correlated with the formation of green pigments under alkaline conditions in the presence of ambient O_2 . However, incubation of soybean *Lba* under similar conditions did not result in the appearance of green pigments (Lehtovaara, 1978). Spectrally, the green pigments obtained from alkaline oxidation of bean *Lb* were quite different from those reported in this study. Soybean *Lbs*, however, contain a conserved Tyr¹³³ residue near the COOH terminus, which may participate in oxidation reactions (Davies and Puppo, 1992). Also, this residue interacts with the heme (Ollis et al., 1983). Oxidation of Tyr¹³³ could lead to perturbation of the heme pocket with resultant changes in ligand binding. Furthermore, any chemical changes in the Tyr ring structure would be expected to affect its absorption in the UV and could thus be detected in the apo-protein. *Lba* contains two Trp and three Tyr residues (Hyldig-Nielsen et al., 1982), and, therefore, the expected ratio of Tyr to Trp is 1.5. A spectroscopic investigation of the Tyr to Trp ratios in apo-*Lba* and apo-*Lbams* revealed no differences for any of the samples analyzed (Table I). The lack of differences would indicate that there was no chemical change to these residues,




Figure 2. Visible absorption spectra of Lb^{2+} and Lb^{2+} nicotinate complexes for *Lba* and *Lbam*s. Conditions of analyses were as described in Figure 1, except *Lb* samples were reduced as described in "Materials and Methods." Lb^{2+} nicotinate complexes were formed by the addition of 0.5 mM (final concentration) of nicotinate to the reduced *Lb* samples.




Figure 3. Visible absorption spectra for $Lb^{2+}O_2$ and $Lb^{2+}CO$ complexes for Lba and Lbams. Conditions were as described in Figures 1 and 2, except reduced Lb samples were incubated in O_2 - or CO -saturated buffer at 22°C for 20 min.

especially to Tyr. These measurements are based on differences in the extinction coefficients as a function of pH (Edelhoc, 1967), and it is conceivable that oxidation of Tyr could have been reversed during sample handling. Attempts to chemically acylate accessible Tyr residues on Lba and Lbams with a variety of reagents (Riordan et al., 1965; Cuatrecasas

et al., 1968) did not yield useful data. Such chemical reactions were not sufficiently specific and resulted in a progressive denaturation of the intact proteins (data not shown).

In addition to Tyr¹³³, Leu¹³⁴ and Ile¹³⁹ near the COOH terminus also interact with the heme of soybean Lb (Ollis et al., 1983); these residues form a hydrophobic end to the heme pocket. The results of earlier studies had indicated that there was no apparent change to the NH₂ terminus of the Lbam proteins (Jun, 1993) and that proteolysis might be involved in the conversion of Lba to Lbam (Wagner and Sarath, 1987). Thus, proteolytic cleavage of residues from the COOH terminus might result in changing the environment of the heme pocket with resultant effects on ligand binding. To determine fully the nature of any of such modifications, IEF-purified Lbs were subjected to trypsin digestion. Peptide profiles resulting from tryptic digestion are shown in Figure 4. The overall profiles and the relative ratios of the peaks were generally similar for all of the proteins analyzed. To determine the residues near the COOH terminus, peptide peaks exhibiting high A_{290} were collected and subjected to automated Edman degradation. The results of this analysis identified the COOH terminal fragments from residues 121 to 127 (peptide labeled B in Fig. 4) and from residues 128 to 141 (peptide labeled A in Fig. 4). Conceivably, deamidation of Asn and Gln residues in Lba could result in changing the isoelectric point of such modified molecules (Jun, 1993). In addition, such changes might have an effect on the physical and chemical characteristics of the deamidated proteins. However, the similar peptide profiles of Lbams and Lba observed upon tryptic digestion (Fig. 4) do not support such a change.

Overall, these results ruled out any gross modification of the protein backbone in the Lbam molecules and suggested that there might be changes to the heme.

Pyridine Hemochromogen Spectra

Pyridine hemochromogen spectra of the heme from Lba and Lbams provided strong evidence for changes to the heme (Fig. 5). As expected, Lba hemochromogen spectra were

Table 1. Spectrophotometric determination of the ratio of Tyr to Trp in lysozyme, sperm whale apomyoglobin, and apoleghemoglobin

Apoproteins were prepared by 2-butanone method (Ascoli et al., 1981) and dissolved in 20 mM K-phosphate buffer (pH 7.0) containing 6 M guanidine chloride at a concentration of 20 mM. The pH 12.0 was obtained by adding 5 μ L of 10 N KOH to the neutral apoprotein solutions. The ratio of Tyr to Trp for these samples was determined using the spectrophotometric procedure of Edelhoc (1967).

	$A_{280, pH7}$	$A_{300, pH7}$	$A_{300, pH12}$	Ratio of Tyr to Trp	
				Calculated	Expected
Lysozyme ^a	0.699	0.149	0.276	0.51	0.5
SWMb ^b	0.320	0.097	0.234	1.42	1.5
Lba ^c	0.316	0.109	0.247	1.45	1.5
Lbam ₁	0.321	0.097	0.237	1.45	
Lbam ₂	0.296	0.105	0.234	1.45	
Lbam ₃	0.299	0.123	0.252	1.44	

^a Lysozyme from egg white contains three Tyr and six Trp. ^b SWMb, sperm whale myoglobin, contains three Tyr and two Trp. ^c Soybean Lba contains three Tyr and two Trp.

identical with those of free hemein with absorption peaks at 556 and 525 nm and with a small peak at 480 nm. Spectra obtained from *Lbam* samples also had absorption peaks at 556 and 525 nm that were of reduced amplitude, none at 480 nm, and a new peak at 580 nm. These spectral differences suggested that the heme in *Lbams* was modified in some way. However, the presence of apparent α and β bands at 556 and 525 nm, respectively, in the *Lbams* indicated that the macroporphyrin ring was intact, and the native spectra indicated that the heme contained iron. Major changes to the porphyrin ring would result in the release of iron as well as a failure to detect the α and β bands at 556 and 525 nm, respectively. Degradation products of heme, such as biliverdin, exhibit a linear pyrrole ring structure and show a broad absorption spectra between 400 and 750 nm with a peak centered at 665 nm (Heirwegh et al., 1991). In animal systems, heme degradation is known to occur by hydroxylation of the methene bridge (Schmid and McDonagh, 1975) with further catabolism of the free heme catalyzed by the enzyme heme oxygenase (Brown et al., 1990). Heme destruction can also be initiated by reductants such as ascorbate in the




Figure 4. Reversed phase HPLC chromatograms of trypsin-digested Lb samples. Lb samples (0.4 mg) were digested initially with an 8- μ g aliquot of *N*-tosyl-L-Phe chloromethyl ketone-treated trypsin in 0.4 mL of 20 mM Tris-HCl (pH 8.0) containing 10 mM CaCl_2 for 3 h at 37°C, followed by a second 8- μ g aliquot of trypsin for additional 2 h. The digestion mixtures were separated by HPLC on a C_{18} - μ Bondapak column (250 \times 4.6 mm), and the eluate was monitored at 215 nm. Peaks labeled A and B were further subjected to automatic Edman degradation. Conditions: solvent A, 0.1% (v/v) TFA; solvent B, 90% (v/v) acetonitrile in 0.1% (v/v) TFA; flow rate, 1 mL min^{-1} . The gradient parameters were 0 to 30 min, 0 to 40% solvent B in a linear gradient; 30 to 35 min, 40 to 100% in a linear gradient; 35 to 40 min, 100% isocratic; 40 to 45 min, 100 to 0% in a linear gradient.




Figure 5. Pyridine hemochromogen spectra of Lb samples. Lb samples (1 nmol) or hemein (0.1 mL) were prepared in 0.3 mL of alkaline pyridine solution composed of 33.3% (v/v) pyridine and 0.1 N NaOH. The sample mixtures (0.4 mL) were reduced with an excess of solid sodium dithionite (a few crystals). Numbers on the graphs indicate wavelength of the peaks.

presence of O_2 (Buldain et al., 1986), by thiol compounds that result in the formation of sulfur-heme adducts (Luiz-Dafré and Reischl, 1990), or by H_2O_2 under physiological conditions (Prasad et al., 1989). Root nodules contain an abundance of ascorbate and reduced thiols such as GSH (Dalton et al., 1991). Ascorbate in the presence of O_2 will convert bean Lbs into inactive forms; however, this reaction requires nonphysiological conditions (Lehtovaara, 1978). The pH of functional root nodules is near neutral and becomes more acidic during senescence (Klucas, 1974; Pladys et al., 1988), suggesting that heme destruction might involve other mechanisms, such as reactions with the hydroxyl radical that is produced during stress (Becana and Klucas, 1992). Our data suggest that functional Lbs can be converted in vivo into proteins exhibiting aberrant binding of O_2 ; however, the nature of this change is unknown.

Lbams appear to have sustained some form of modification to their heme. These hemoproteins do not bind O_2 with characteristics similar to *Lba* and may be functionally impaired. Since these molecules have been shown to become predominant during aging or stress-induced senescence of soybean root nodules, it would appear that *Lbams* are generated in vivo from *Lba*. Thus, *Lbams* could represent an early degradation product of *Lba* and might be used as a marker for the physiological status of the root nodule.

LITERATURE CITED

- Appleby CA (1984) Leghemoglobin and Rhizobium respiration. *Annu Rev Plant Physiol* 35: 433-478
- Appleby CA, Nicola NA, Hurrell JGR, Leach SJ (1975) Characterization and improved separation of soybean leghemoglobins. *Biochemistry* 14: 4444-4450
- Ascoli F, Fanelli MRR, Antonini E (1981) Preparation and properties of apohemoglobin and reconstituted hemoglobins. *Methods Enzymol* 76: 72-87
- Becana M, Klucas RV (1990) Enzymatic and nonenzymatic mechanisms for ferric leghemoglobin reduction in legume root nodules. *Proc Natl Acad Sci USA* 87: 7295-7299
- Becana M, Klucas RV (1992) Transition metals in legume root nodules: iron-dependent free radical production increases during nodule senescence. *Proc Natl Acad Sci USA* 89: 8958-8962
- Becana M, Salin ML, Ji L, Klucas RV (1991) Flavin-mediated reduction of ferric leghemoglobin from soybean nodules. *Planta* 183: 575-583
- Brown SB, Houghton JD, Wilks A (1990) Heme degradation and biosynthesis of bilins. In HA Dailey, ed, *Biosynthesis of Heme and Chlorophylls*. McGraw-Hill, New York, pp 543-575
- Buldain G, Frydman RB, Tomaro ML, Frydman B (1986) The chemical and enzymatic oxidation of hematoheemin IX: identification of hematobiliverdin IXa as the sole product of the enzymatic oxidation. *Eur J Biochem* 156: 176-184
- Cuatrecasas P, Fuchs S, Anfinsen CB (1968) The tyrosyl residues at the active site of staphylococcal nuclease. *J Biol Chem* 243: 4787-4798
- Dalton DA, Post CJ, Langeberg L (1991) Effect of ambient oxygen and of fixed nitrogen on concentrations of glutathione, ascorbate, and associated enzymes in soybean root nodules. *Plant Physiol* 96: 812-818
- Davies MJ, Puppo A (1992) Direct detection of globin-derived radical in leghemoglobin treated with peroxides. *Biochem J* 281: 197-201
- Edelhoc H (1967) Spectroscopic determination of tryptophan and tyrosine in proteins. *Biochemistry* 6: 1948-1954
- Heirwegh KP, Blanckaert N, Van Hees G (1991) Synthesis, chromatographic purification, and analysis of isomers of biliverdin IX and bilirubin IX. *Anal Biochem* 195: 273-278
- Hyldig-Nielsen JJ, Jensen EO, Paludan K, Wiborg O, Garrett R, Jorgensen P, Marcker EA (1982) The primary structure of two leghemoglobin genes from soybean. *Nucleic Acids Res* 10: 689-701
- Jun H-K (1993) Isolation and characterization of modified forms of soybean leghemoglobin a. PhD thesis, University of Nebraska-Lincoln
- Klucas RV (1974) Studies on soybean nodule senescence. *Plant Physiol* 54: 612-616
- Lehtovaara P (1978) Oxidation of glycine by *Phaseolus* leghemoglobin with associate catabolic reactions at the haem. *Biochem J* 176: 351-358
- Luiz-Dafré A, Reischl E (1990) High hemoglobin mixed disulfide content in hemilysates from stress shark. *Comp Biochem Physiol* 96B: 215-219
- Ollis DL, Appleby CA, Colman PM, Cutten AE, Guss JM, Venkatappa MP, Freeman HC (1983) Crystal structure of soybean ferric leghemoglobin a nicotinate at a resolution of 3.3 Å. *Aust J Chem* 36: 451-468
- Pfeiffer NE, Malik NSA, Wagner FW (1983) Reversible dark-induced senescence of soybean root nodules. *Plant Physiol* 71: 393-399
- Pladys D, Barthe P, Rigaud J (1988) Changes in intracellular pH in French bean nodules induced by senescence and nitrate treatment. *Plant Sci* 56: 99-106
- Prasad MR, Engelman RM, Jones RM, Das DK (1989) Effects of oxyradicals on oxymyoglobin: deoxygenation, haem removal and iron release. *Biochem J* 263: 731-736
- Riggs A (1981) Preparation of blood hemoglobins of vertebrates. *Methods Enzymol* 76: 5-29
- Riordan JF, Wacker WEC, Vallee BL (1965) N-Acetylimidazole: a reagent for determination of "free" tyrosyl residues of proteins. *Biochemistry* 4: 1758-1765
- Sarath G, Cohen HC, Wagner FW (1986) High performance liquid chromatographic separation of leghemoglobins from soybean root nodules. *Anal Biochem* 154: 224-231
- Schmid R, McDonagh AF (1975) The enzymatic formation of bilirubin. *Ann NY Acad Sci* 244: 533-552
- Sutton WD (1983) Nodule development and senescence. In WJ Broughton, ed, *Nitrogen Fixation*, Vol 3. Clarendon Press, Oxford, UK, pp 144-212
- Wagner FW, Sarath G (1987) Biochemical changes in stressed and senescent soybean root nodules. In ED Thompson, NA Nothnagel, RC Huffaker, eds, *Plant Senescence: Its Biochemistry and Physiology*. The American Society of Plant Physiologists, Rockville, MD, pp 190-197